



LIST OF REFERENCES CITED BY APPLICANT

(Use several sheets if necessary)

ATTY. DOCKET NO.

10624-048-999

APPLICATION NO.

09/385,918

APPLICANT

Hoekstra et al.

FILING DATE

October 30, 1999

GROUP

1646

U.S. PATENT DOCUMENTS

*EXAMINER INITIAL	DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE

FOREIGN PATENT DOCUMENTS

DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION
					YES NO
AA	WO 99/01765	1/14/99	PCT		

OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.)

AB	Attisano et al., 1993, "Identification of human activin and TGF beta type I receptors that form heteromeric kinase complexes with type II receptors", Cell 75(4):671-80.
AC	Attisano et al., 1996, "Activation of signalling by the activin receptor complex", Mol Cell Biol. 16(3):1066-73.
AD	Cui et al., 1996, "TGFbeta1 inhibits the formation of benign skin tumors, but enhances progression to invasive spindle carcinomas in transgenic mice", Cell 86(4):531-42.
AE	Derynck, 1998, "Smads: transcriptional activators of TGF-beta responses", Cell 95(6):737-40. R
AF	Hayashi et al., 1997, "The MAD-related protein Smad7 associates with the TGFbeta receptor and functions as an antagonist of TGFbeta signaling", Cell 89(7):1165-73.
AG	Heldin et al., 1997, "TGF-beta signalling from cell membrane to nucleus through SMAD proteins", Nature 390: 465-471.
AH	Hershko et al., 1983, "Components of ubiquitin-protein ligase system. Resolution, affinity purification, and role in protein breakdown", J Biol Chem. 258(13):8206-14.
AI	Hojo et al., 1999, "Cyclosporine induces cancer progression by a cell-autonomous mechanism", Nature 397(6719):530-4.
AJ	Huibregtse et al., 1995, "A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase. Proc Natl Acad Sci U S A. 92(7):2563-7.
AK	Huibregtse et al., 1993, "Cloning and expression of the cDNA for E6-AP, a protein that mediates the interaction of the human papillomavirus E6 oncoprotein with p53", Mol Cell Biol. 13(2):775-84.
AL	Imamura, 1997, "Smad6 inhibits signalling by the TGF-beta superfamily", Nature 389(6651):622-6
AM	Kretschmar et al., 1997, "Opposing BMP and EGF signalling pathways converge on the TGF-beta family mediator Smad1", Nature 389(6651):618-22.
AN	Kretschmar and Massague, 1998, "SMADS: mediators and regulators of TGF-beta signaling", Curr Opin Genet Dev. 8(1):103-11
AO	Lin et al., 1992, "Expression cloning of the TGF-beta type II receptor, a functional transmembrane serine/threonine kinase", Cell 68(4):775-85.
AP	Liu et al., 1996, "A human Mad protein acting as a BMP-regulated transcriptional activator", Nature 381(6583):620-3.
AQ	Markowitz et al., 1995, "Inactivation of the type II TGF-beta receptor in colon cancer cells with microsatellite instability", Science 268(5215):1336-8.
AR	Mathews and Vale, 1991, "Expression cloning of an activin receptor, a predicted transmembrane serine kinase", Cell. 65(6):973-82.
AS	Miyazono et al., 1999, "Signal transduction by bone morphogenetic protein receptors", Bone 25(1): 91-93.
AT	Nakao et al., 1997, "Identification of Smad2, a human Mad-related protein in the transforming growth factor beta signaling pathway", J Biol Chem. 272(5):2896-900.
AU	Newfeld et al., 1996, "Mothers against dpp encodes a conserved cytoplasmic protein required in DPP/TGF-beta responsive cells", Development 122(7):2099-108.

Am	AV	Pirozzi et al., 1997, "Identification of novel human WW domain-containing proteins by cloning of ligand targets", J Biol Chem. 272(23):14611-6.
	AW	Reynisdottir et al., 1995, "Kip/Cip and Ink4 Cdk inhibitors cooperate to induce cell cycle arrest in response to TGF-beta. Genes Dev. 9(15):1831-45.
	AX	Sekelsky et al., 1995, "Genetic characterization and cloning of mothers against dpp, a gene required for decapentaplegic function in Drosophila melanogaster", Genetics 139(3):1347-58.
	AY	Staub et al., 1996, " WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na ⁺ channel deleted in Liddle's syndrome", EMBO J. 15(10):2371-80.
	AZ	Sudol et al., 1996, "Structure and function of the WW domain", Prog Biophys Mol Biol. 5(1-2):113-32
	BA	Tsuchida et al., 1993, "Cloning and characterization of a transmembrane serine kinase that acts as an activin type I receptor", Proc Natl Acad Sci U S A. 90(23):11242-6.
	BB	Ulloa et al., 1999, "Inhibition of transforming growth factor-beta/SMAD signalling by the interferon-gamma/STAT pathway. Nature 397(6721):710-3.
	BC	Zawel et al., 1998, "Human Smad3 and Smad4 are sequence-specific transcription activators", Mol Cell. 1(4):611-7.
	BD	Zhang et al., 1996, "Receptor-associated Mad homologues synergize as effectors of the TGF-beta response", Nature 383(6596):168-72.
	BE	Zhu et al., 1998, "Smad3 mutant mice develop metastatic colorectal cancer", Cell 94(6):703-14.
↓	BF	Zhu et al., 1999, "A SMAD ubiquitin ligase targets the BMP pathway and affects embryonic pattern formation", Nature 400: 687-93.
EXAMINER <i>J. H. Ch</i>		DATE CONSIDERED 07/17/01
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.		